

[All Databases](#)[PubMed](#)[Nucleotide](#)[Protein](#)[Genome](#)[Structure](#)[OMIM](#)[PMC](#)[Journals](#)[Books](#)

Search PubMed

for

Go

Clear

[Limits](#) [Preview/Index](#) [History](#) [Clipboard](#) [Details](#)

Display Abstract

Show

20

Sort by

Send to

[About Entrez](#)[Text Version](#)[Entrez PubMed](#)[Overview](#)[Help | FAQ](#)[Tutorial](#)[New/Noteworthy](#)[E-Utilities](#)[PubMed Services](#)[Journals Database](#)[MeSH Database](#)[Single Citation Matcher](#)[Batch Citation Matcher](#)[Clinical Queries](#)[Special Queries](#)[LinkOut](#)[My NCBI \(Cubby\)](#)[Related Resources](#)[Order Documents](#)[NLM Mobile](#)[NLM Catalog](#)[NLM Gateway](#)[TOXNET](#)[Consumer Health](#)[Clinical Alerts](#)[ClinicalTrials.gov](#)[PubMed Central](#)

1: Virology. 1988 Nov;167(1):8-14.

[Related Articles, Links](#)

Amino acid substitution at position 226 of the hemagglutinin molecule of influenza (H1N1) virus affects receptor binding activity but not fusion activity.

Nobusawa E, Nakajima K.

Institute of Medical Science, University of Tokyo, Japan.

The receptor binding site of the hemagglutinin (HA) molecule of type A influenza virus A/USSR/90/77 (H1N1) has been studied. Site-specific mutagenesis has been used to introduce base changes into the sequence that codes for the amino acid residue at position 226 on the HA molecule, and mutant sequences replaced the wild-type sequence of the HA gene of the SV40-HA recombinant virus (SVHA). Mutant HA proteins were expressed in African green monkey kidney cells and analyzed for receptor binding and fusion activities. Two mutant HA proteins containing single amino acid substitutions of Asn and Met for Gln at position 226 retained their receptor binding activity, but others with amino acid substitutions Glu, His, Leu, Val, and Thr for Gln at position 226 lost this activity. All the mutant proteins retained their fusion activity. On the other hand, another four mutants containing single amino acid substitutions at positions other than 226 retained the receptor binding and fusion activities, despite the drastic change in charge or polarity to the respective amino acids. These results suggest that amino acid residue 226 of the H1 subtype of HA is critical for receptor binding activity of the HA protein. Our results also show that the lack of receptor binding activity of the HA protein does not affect fusion activity.

PMID: 2460997 [PubMed - indexed for MEDLINE]

Display Abstract



Show

20



Sort by



Send to

[Write to the Help Desk](#)[NCBI | NLM | NIH](#)[Department of Health & Human Services](#)[Privacy Statement | Freedom of Information Act | Disclaimer](#)

